

Comparison of the thin flux tube approximation with 3D MHD simulations

L. Yelles Chaouche^{1,2}, S. K. Solanki^{1,3}, and M. Schüssler¹

¹ Max-Planck-Institut für Sonnensystemforschung, Max-Planck-Strasse 2, 37191 Katlenburg-Lindau, Germany

² Instituto de Astrofísica de Canarias, C/ Vía Láctea, s/n, E38205 - La Laguna (Tenerife). España

³ School of Space Research, Kyung Hee University, Yongin, Gyeonggi 446-701, Korea
e-mail: yelles@mps.mpg.de

Preprint online version: October 9, 2009

ABSTRACT

Context. The structure and dynamics of small vertical photospheric magnetic flux concentrations has been often treated in the framework of an approximation based upon a low-order truncation of the Taylor expansions of all quantities in the horizontal direction, together with the assumption of instantaneous total pressure balance at the boundary to the non-magnetic external medium. Formally, such an approximation is justified if the diameter of the structure (a flux tube or a flux sheet) is small compared to all other relevant length scales (scale height, radius of curvature, wavelength, etc.). The advent of realistic 3D radiative MHD simulations opens the possibility of checking the consistency of the approximation with the properties of the flux concentrations that form in the course of a simulation.

Aims. We carry out a comparative analysis between the thin flux tube/sheet models and flux concentrations formed in a 3D radiation-MHD simulation.

Methods. We compare the distribution of the vertical and horizontal components of the magnetic field in a 3D MHD simulation with the field distribution in the case of the thin flux tube/sheet approximation. We also consider the total (gas plus magnetic) pressure in the MHD simulation box.

Results. Flux concentrations with super-equipartition fields are reasonably well reproduced by the second-order thin flux tube/sheet approximation. The differences between approximation and simulation are due to the asymmetry and the dynamics of the simulated structures.

Key words. Magnetohydrodynamics (MHD) – Sun: magnetic fields – Sun: photosphere

1. Introduction

Much of the solar photospheric magnetic flux exists in the form of discrete concentrations in intergranular lanes having a field strength of 1-2 kG (Stenflo 1973; Wiehr 1978; Rüedi et al.

1992; Rabin 1992; Martínez Pillet et al. 1997), for reviews see Solanki (1993); Solanki et al. (2006).

Theoretical models of these flux concentrations have widely used the concept of the flux tube: a bundle of field lines with circular cross-section separated from the non-magnetic environment by a tangential discontinuity (see e.g. Schüssler 1992). Such a structure can be described, under certain conditions, by the so called "thin flux tube approximation". In its simplest form, the axial component of the magnetic field is assumed to be constant across the tube's cross-section, while the radial component is a linear function of the radial coordinate (Defouw 1976; Roberts & Webb 1978, 1979). The thin flux tube approximation can be formally justified if the diameter of the flux tube is sufficiently small compared to variations of the relevant physical quantities (such as pressure, density,...etc.) along the tube's cross-section (Spruit 1981; Schüssler 1992).

The equations describing a thin flux tube can be obtained by writing all physical quantities (magnetic field, temperature, pressure,...etc) in terms of a Taylor expansion in the radial distance from the axis, and inserting them in the MHD equations. By collecting terms of similar order one obtains a hierarchy of equations (Ferriz-Mas & Schüssler 1989). Truncating this hierarchy after the 1st order allows the 0th-order approximation introduced above to be obtained.

Extensions of the thin flux tube approximation to higher orders have been given in the literature. By retaining second-order terms Pneuman et al. (1986) have included in their modelling the effects of field line curvature, internal structures, twist, and the merging of flux tubes with their neighbours. A derivation of linear wave modes of a flux tube up to second order has been carried out by Ferriz-Mas et al. (1989).

There is a large body of work in the literature based upon the thin flux tube approximation. This includes theoretical work (structure of flux concentrations, equilibrium, oscillations/wave, stability,...etc.) and interpretation of observations (for reviews, see Solanki 1993; Solanki et al. 2006). Various aspects of the thin tube approximation have been compared to observational data (e.g. Zayer et al. 1989; Bruls & Solanki 1995; Solanki et al. 1996), but its validity has not been tested on the basis of the most advanced numerical simulations.

In the last two decades, the possibilities to self consistently model magneto-convection at the solar photosphere using the full set of MHD equations including radiative and convective energy transport (e.g. Nordlund 1983; Stein & Nordlund 1998; Bercik 2002; Stein & Nordlund 2003; Vögler & Schüssler 2003; Vögler et al. 2005) have greatly improved. The structure of flux concentrations in such MHD simulations appears rather complex, owing to their interaction with convection and energy exchange with the neighbouring plasma.

We aim to evaluate to which extent the magnetic structures forming in 3D MHD simulations can be described using the thin flux tube/sheet approximation.

2. A series expansion of the thin flux tube/sheet equations

We consider a magnetic flux tube to be a bundle of magnetic field lines with a circular cross section, which is separated from its non-magnetic surroundings by a tangential discontinuity with a surface current. For an axisymmetric vertical flux tube, we adopt cylindrical coordinates (r, θ, z) , with the z -axis pointing in the vertical direction. Physical quantities are regular at the axis ($r = 0$), so that they can be described in terms of a Taylor expansion in the radial coordinate (Roberts & Webb 1978; Spruit 1981; Pneuman et al. 1986; Ferriz-Mas et al. 1989; Ferriz-Mas & Schüssler 1989)

The properties of the axisymmetric MHD equations (Ferriz-Mas & Schüssler 1989) imply that only even orders are non-zero in the above-mentioned expansions for scalar quantities (such as temperature or density) and for z -components of vectors, whereas for the radial and θ -components of vectors only the odd orders remain.

The three components of the magnetic field vector, the temperature and the pressure can be written in a non-dimensional way:

$$b_z = h_0 + h_2 x^2 + h_4 x^4 + \dots, \quad (1)$$

$$b_r = f_1 x + f_3 x^3 + f_5 x^5 + \dots, \quad (2)$$

$$b_\theta = s_1 x + s_3 x^3 + s_5 x^5 + \dots \quad (3)$$

$$p = p_0 + p_2 x^2 + p_4 x^4 + \dots \quad (4)$$

$$\sigma = \sigma_0 + \sigma_2 x^2 + \sigma_4 x^4 + \dots, \quad (5)$$

with $p = P/P^*$, $\sigma = T/T^*$, $b = B/B^*$, $x = r/H^*$, $y = z/H^*$, $H^* = kT^*/(m_p g)$ and $\alpha = 4\pi P^*/B^{*2}$. Where B_r , B_θ , B_z , represent the three components of the magnetic field vector. P and T are the gas pressure and temperature, respectively. The quantities with an asterisk are defined at the tube's axis ($x = r = 0$) and at a reference height ($z = y = 0$). k is Boltzmann's constant, m_p the mean particle mass, g the gravitational acceleration, and H the scale height.

2.1. B_z and B_r under the thin flux tube approximation

Following, e.g., Pneuman et al. (1986), in a static atmosphere, we insert the expansions (1 to 5) in the three components of the momentum equation and the solenoidality relation, and collect terms of equal power in x into equations of corresponding order. Considering equations including terms up to the third order, and assuming that the flux tubes studied here have negligible twist, we obtain the following relations (Pneuman et al. 1986):

$$h_2 = -\frac{1}{4}h_0'' - \frac{\alpha p_2}{h_0}, \quad (6)$$

$$f_1 = -\frac{1}{2}h_0', \quad (7)$$

and

$$f_3 = -\frac{1}{4}h'_2, \quad (8)$$

where the prime indicates a derivative with respect to y

We can then deduce B_z up to the second order, and B_r up to the third order.

In order to close the above system it is necessary to consider relations expressing magnetic flux conservation through the tube's cross-section and total pressure balance at the boundary of the flux tube at any height (Ferriz-Mas & Schüssler 1989; Ferriz-Mas et al. 1989). In addition to these relations, Ferriz-Mas et al. (1989) have considered an energy equation, whereas Pneuman et al. (1986) have chosen to prescribe two quantities, such as σ_0 and σ_2 , which allows more flexibility in defining the atmosphere. In order to construct a thin flux tube which we will compare with flux concentrations in MHD simulations, we take h_0 and p_2 from the MHD simulations. Then B_z and B_r are determined from equations (6) to (8). The cross section of the flux tube is determined through the magnetic flux conservation relation:

$$\int_0^{x_0} (h_0 + h_2 x^2) dx = \text{Flux at base of tube} = \text{constant}, \quad (9)$$

where x_0 is the tube's radius at a given height. The total pressure balance can be expressed as:

$$(\beta p_i + b_i^2)|_{x=x_0} = (8\pi P_e + B_e^2)/B_e^{*2}|_{x=x_0}, \quad (10)$$

where $\beta = 8\pi P^*/B^{*2}$, the suffixes i and e indicate internal and external quantities respectively and capital letters indicate dimensional quantities.

Under the 0^{th} -order approximation Eq. (10) reduces to:

$$(\beta p_0 + h_0^2)|_{x=x_0} = (8\pi P_e + B_e^2)/B_e^{*2}|_{x=x_0}, \quad (11)$$

This relation does not depend on the radius of the flux tube. Thus the total pressure at a given height under the 0^{th} -order approximation is constant across the tube's cross-section.

Under the 2^{nd} -order approximation we get:

$$(\beta p_0 + h_0^2) + x^2(\beta p_2 + f_1^2 + 2h_0 h_2)|_{x=x_0} = (8\pi P_e + B_e^2)/B_e^{*2}|_{x=x_0}, \quad (12)$$

In this case the total pressure varies inside the flux tube, but has to match the external total pressure at the tube's boundary.

2.2. B_z and B_x under the thin flux sheet approximation

A flux sheet is an elongated structure with a small width (which we refer to as "W") compared to its length ("L") along the solar surface, i.e. $W \ll L$ at the solar surface. A similar approach as described in the previous section can be used to describe a thin flux sheet. In this case the magnetic field component parallel to L is constant, and thus plays no direct role in the hydrostatic equilibrium. We can then adopt a Cartesian 2D geometry in the $x-z$ plane, where z is the vertical coordinate and x is the horizontal coordinate perpendicular to the vector L .

In a similar way to Sect. 2.1, we can determine h_2 , f_1 and f_3 as functions of h_0 and p_2 .

$$h_2 = -\frac{1}{2}h_0'' - \frac{\alpha p_2}{h_0} \quad (13)$$

$$f_1 = -h_0' \quad (14)$$

and

$$f_3 = -\frac{1}{3}h_2' \quad (15)$$

Note the similarity between these equations and the ones describing the thin flux tube. The main difference (apart of the geometry) is the numerical values of the constant coefficients which affects, for instance, the expansion rate of the flux tube/sheet with height.

3. The radiative MHD simulations

Three dimensional radiation-MHD simulations of the solar photosphere have been described by (Nordlund 1983; Nordlund & Stein 1990; Stein & Nordlund 1998; Bercik 2002; Stein & Nordlund 2003; Vögler & Schüssler 2003; Vögler et al. 2005).

The simulation run used here has been obtained with the fully compressible MURaM code (Vögler 2003; Vögler et al. 2005). It takes into account non-local and non-gray radiative energy transport, and includes the effects of partial ionization. The simulation box has a horizontal extension of $6 \times 6 \text{ Mm}^2$ and is 1.4 Mm deep. The grid cell size is 5 km in the horizontal direction and 7 km in the vertical. The simulation run starts from a plane-parallel atmosphere which extends from -0.8 Mm below to 0.6 Mm above a reference 0, which is roughly situated -100 km below the average continuum optical depth unity ($\tau_{5000} = 1$, which corresponds to the solar surface at 5000 \AA). After convection has fully developed, a mixed-polarity magnetic field configuration with zero net vertical flux is introduced. This is done such that the simulation domain is divided into four parts with vertical field of alternate polarities in a chessboard pattern. We choose a representative snapshot for our analysis (see Fig. 1). The mean unsigned field strength at optical depth unity is 200 G for this snapshot.

4. Analysis of the total pressure in the whole simulation domain

Figure 2 shows horizontally averaged gas and total pressures as a function of height. The solid line represents the gas pressure averaged over regions with field strength smaller than 50 G. The dash-dotted line indicates the gas pressure averaged over magnetic regions. The threshold in $|B_z|$ defining magnetic regions varies linearly from 500 G at the bottom of the simulation box to 300 G at the top. The dashed line represents the total pressure ($P_{\text{tot}} = P + B^2/(8\pi)$) averaged over magnetic regions. The plasma $\beta = 8\pi P/B^2$ for magnetic regions is indicated by square symbols.

The difference between the gas pressures inside and outside magnetic regions becomes smaller with depth. This is due to the large values of the plasma β in the deep layers (e.g. below -400 km) which indicate that the pressure balance between magnetic features and their




Fig. 1. Upper panel: vertical component of the magnetic field (B_z) at a fixed geometrical height near the averaged visible solar surface ($\tau_{5000} = 1$). Lower panel: vertical component of the velocity (V_z) at $\tau_{5000} = 1$. Downflows are represented in red and upflows in blue. The black contours outline regions where $|B_z| > 500$ G. The black vertical lines indicate locations where we carry out a detailed analysis of magnetic elements in Section 5.

surroundings is mainly ensured by gas pressure. Above 300 km, the total pressure in magnetic features shows an excess compared to gas pressure in nearly field-free areas. This excess increases with height and is due to the effect of curvature forces. This implies that the 0^{th} -order thin flux tube/sheet approximation is not sufficient to describe the flux concentrations in the upper part of the simulation box. The gas pressure in nearly field-free regions is higher than the total pressure in the flux concentrations in the height-range situated between 50 and 300 km. This slight pressure excess mainly results from the fact that the gas pressure at equal geometrical height is, on average, higher in the granular upflows than in the intergranular downflow lanes, where the magnetic flux concentrations reside. In addition, a pressure deficit in the flux concentrations relative to their local environment could arise as the result of the outward curvature force




Fig. 2. Horizontal average of the plasma β , gas pressure and total pressure as a function of height. Plotted are, the logarithmic values of the gas pressure averaged over regions with field strength smaller than 50 G (full line), the gas pressure averaged in magnetic flux concentrations (dash-dotted line), the total pressure over magnetic flux concentrations (dashed line) and the plasma β (squares; scale on the right).

of the expanding tubes between 0 and 300 km height. Above 300 km, the sign of the curvature force is reversed as a result of the wineglass shape of the flux tubes caused by the presence of neighboring tubes (reflected in our simulation by the vertical-field upper boundary condition). In any case, the deviation from total pressure balance is very small below 300 km height.

The total pressure balance between a magnetic flux concentration and its non-magnetic surroundings results from the continuity of the normal stress at the boundary separating the flux concentration from its surroundings. In the 0^{th} -order approximation (Eq 11), P_{tot} not only matches the boundary value but is also constant across the flux concentration. The presence of 2^{nd} -order terms (or higher-orders) produces higher or lower values of P_{tot} at the center of flux concentrations (Eq 12) in comparison to P_{tot} at the magnetic/non-magnetic boundary, which remains equal to the external pressure. Thus the total pressure can be used as diagnostic for determining whether a flux concentration has 0^{th} or higher-order configuration. In order to illustrate the distributions of pressures and magnetic field in the simulation box, which includes different sizes and shapes of magnetic concentrations, we show in Figure 3 maps of the gas and total pressures as well as B_z at three heights -98 , 182 and 462 km, where the reference height 0 is roughly situated -100 km below the average continuum optical depth unity, $\tau_{5000} = 1$.




Fig. 3. Gas pressure, total pressure and vertical component of the magnetic field at three geometrical heights -98 , 182 and 462 km. The reference height "0" is located at -100 km below the average continuum optical depth unity at 5000 \AA . A common color table is used for the three B_z maps. At each height, the pressures share the same color table indicated on the right-side of the maps. The black contours enclose regions where $|B_z|$ is higher than 500 G at -98 km, higher than 400 G at 182 km and higher than 300 G at 462 km.

The vertical component of the magnetic field at -98 km is displayed in the lower left panel of Figure 3. Note that the flux resides mainly in strong flux concentrations located in intergranular lanes. The middle panel of the lower row in Figure 3 represents gas pressure at -98 km. Locations where the magnetic flux density is higher than 500 G are outlined by dark contours. The gas pressure is higher at centers of granules comparing to intergranules. This pressure excess drives the horizontal flows towards the intergranular lanes (see e.g. Stein & Nordlund 2003). Intergranular lanes display a mixed picture with high gas pressure (which stops the horizontal flows) but also low pressure areas. The magnetic flux concentrations show lower gas pressure.

The total pressure inside flux concentrations at -98 km (lower right panel of Figure 3) is roughly close to gas pressure outside, and does not vary significantly within individual flux concentrations. Constant P_{tot} is a necessary condition (but not sufficient) for the validity of the 0^{th} -order thin flux tube/sheet model.

The existence of 2^{nd} -order terms (or higher-orders) in pressure and magnetic field leads to higher/lower values of the total pressure at the center of flux concentrations. So, one way of measuring the importance of higher-order terms is to compute the standard deviation and the mean value of the total pressure inside magnetic elements and compare them with the corresponding values outside magnetic regions (see Table 1).

Table 1. Standard deviations and mean value of P_{tot}

Altitude [km]	-98	182	462
Standard deviation of P_{tot} in non-magnetic regions (σ_{NM}) [cgs]	27703.0	9707.59	1055.04
Standard deviation of P_{tot} in magnetic regions (σ_M) [cgs]	32709.4	7950.80	2151.41
Mean value of P_{tot} in non-magnetic regions (\overline{NM}) [cgs]	254220.0	57750.9	4901.33
Mean value of P_{tot} in magnetic regions (\overline{M}) [cgs]	249696.0	47409.8	8626.20
$\sigma_{NM}/\overline{NM}$	0.108	0.168	0.215
σ_M/\overline{M}	0.130	0.167	0.249

At an altitude of -98 km (Table 1), \overline{M} is slightly lower than \overline{NM} because magnetic flux concentrations are located in intergranular lanes where the pressure at this altitude is slightly lower than the average pressure over the simulation domain. σ_M/\overline{M} is larger than $\sigma_{NM}/\overline{NM}$, this does not result from higher-order terms, but rather indicates the presence of fluctuations inside magnetic elements. This is due to the fact that the plasma beta at this altitude is larger than unity (See Fig. 2) which indicates that convection affects and perturbs the field's regularity. Note that locations with particularly low total pressure (e.g. the green-colored ones) are generally unrelated to magnetic flux concentrations.

At a higher altitude (182 km) we see in Figure 3 that magnetic structures have expanded. The gas pressure has, on average, lower values in intergranular lanes and particularly low values inside magnetic elements. The total pressure is lower in intergranular lanes even when there is no (or low) magnetic field, e.g. in the region around the coordinates (3Mm, 2.8Mm). The mean value \overline{NM} is higher than \overline{M} (Table 1). The normalized fluctuations of P_{tot} inside and outside magnetic elements are similar ($\sigma_M/\overline{M} \approx \sigma_{NM}/\overline{NM}$). Hence there is little evidence for a significant contribution from higher-order terms.

Near the top of the box, at a geometrical height of 462 km, we notice that the total pressure (Figure 3) increases towards the center of flux concentrations. $\overline{M} > \overline{NM}$ and $\sigma_M/\overline{M} > \sigma_{NM}/\overline{NM}$. This indicates that the total pressure is not a 0^{th} -order function. This effect is more pronounced in large flux concentrations. The plasma β is small at these heights (see Fig. 2), thus we expect a nearly force-free equilibrium with a balance between curvature force and magnetic pressure gradient. So the outward magnetic pressure force will be balanced by the inward curvature force.

Thus the magnetic pressure (\approx total pressure) has to increase inward. Hence the increase of P_{tot} at the center of flux concentrations in the upper right panel of Figure 3.

5. Analysis of individual magnetic structures

The flat profiles of total pressure in the lower part of the atmosphere are in favour of the applicability of the 0^{th} -order thin flux tube/sheet approximation. In the upper part of the atmosphere, however, the magnetic features show a total pressure excess in their center. This indicates that the 0^{th} -order thin flux tube/sheet approximation is not applicable, but possibly the extension of the approximation to 2^{nd} -order is sufficient to describe the force equilibrium of the magnetic structures. For a quantitative investigation we select three flux concentrations in the MHD simulation run according to their width and morphology. These will be treated in the next three sub-sections.

5.1. Thin flux sheet

We compare the properties of a narrow flux sheet in the MHD simulation (Fig. 1) with the thin flux sheet model presented in Sect. 2.2. Note that the flux tubes/sheets in a magneto-convection simulation are not static (unlike the assumption made in Sect. 2). They interact with the external plasma, and get distorted by the granulation motion. They also exchange energy (mainly by radiation) with the surroundings. In order to maintain the numerical stability of the simulation, the gradient of any physical quantity cannot be too large between two neighbouring grid cells. More specifically, the magnetic flux density must not jump abruptly from the boundary of a flux tube to the neighbouring non-magnetized plasma (Vögler 2003). Thus the boundary layer separating a flux tube from the surrounding non-magnetized plasma is a few grid points wide, unlike the tangential discontinuity in the case of an ideal flux tube. We wish to see whether simulations and thin flux sheet/tube approximation are consistent with each other in spite of the fact that MHD flux tubes/sheets have finite boundary layers, internal and external dynamics and deviate from an axi- or translationally symmetric configuration.

We select a rather narrow sheet-like structure in the simulation domain. A vertical 2D cut through the flux sheet (shown in Fig. 4) at the location indicated by the dark line in the upper left corner of Fig. 1 reveals the morphology of the magnetic field. The expansion of the flux sheet with height is mainly determined by magnetic flux conservation with height and a horizontal balance between the magnetic plus gas pressure inside the sheet with the gas pressure outside.

Figure 5 shows profiles of gas pressure (full lines) and total pressure (triangles) along the 5 horizontal lines in Figure 4. The location of the magnetic flux concentration is reflected by the lower gas pressure. The vertical dashed lines outline regions where B_z is higher than 0.75 of its maximum value. The profiles indicate that the flux sheet's equilibrium in the lower panels is consistent with balance of total pressure in the zeroth-order thin flux sheet approximation (see Eq 11, which is valid for both flux tubes and flux sheets). In the top panel we see that the total pressure increases somewhat towards the center of the sheet, which indicates the necessity of




Fig. 4. Vertical 2D cut through the MHD simulation box at the location shown by the black line at the upper-left corner in Figure 1. The horizontal lines indicate locations at which we plot profiles of various physical quantities in Figs. 5 to 7.

extending the approximation to second- (or higher-) order (see Eq 12). At this height, the plasma β has become so small that the internal equilibrium becomes nearly force free, i.e., curvature forces and magnetic pressure gradient balance each other.

Figure 6 shows the vertical component of the magnetic field along the 5 horizontal lines in Figure 4. The triangles represent B_z in the case of a thin flux sheet in the second-order approximation. The solid curves represent B_z from the MHD simulations. We note that B_z for the thin flux sheet at the two lower panels is close to a constant (small contribution from the 2^{nd} -order terms), whereas in the three upper panels the second-order terms become more important.

The 2^{nd} -order approximation reproduces reasonably well the overall B_z profiles obtained from the MHD simulations in the higher layers of the atmosphere. The profiles of B_z from the




Fig. 5. Gas pressure (solid lines) and total pressure (triangles) along the 5 horizontal lines in Figure 4. The vertical dashed lines outline regions where B_z is larger than 75 percent of its maximum value.

simulation exhibit some structures across the sheet's cross-section which are not reproduced by the thin sheet model. This is because this latter model produces only symmetric profiles of B_z (Ferriz-Mas & Schüssler 1989). The actual profiles of B_z are asymmetric primarily in the sense that the left part exhibits larger values than the right part. This is associated with lower values of the pressure at these locations, so B_z has to increase in order to keep P_{tot} balanced (see Figure 5).

The distribution of the horizontal field component and its approximation with the thin flux sheet model are shown in Fig. 7. Here B_x includes a third-order term (see section 2.2). The profiles of the actual field are smooth for the three upper panels (low β). In the two lowest panel we notice some fluctuations mainly due to perturbations by the external convection. The fit between the simulation result and the thin sheet model is relatively good for the three upper

Table 2. Relative importance of the average values of the series expansion $\overline{|B2|}/\overline{|B0|}$, $\overline{|B4|}/\overline{|B0|}$ and $\overline{|B3|}/\overline{|B1|}$.

Thin flux sheet: height from reference [km]	-98	42	182	322	462
$\overline{ B2 }/\overline{ B0 }$	0.009	0.014	0.036	0.118	0.106
$\overline{ B4 }/\overline{ B0 }$	7.67 e-05	2.75 e-05	0.001	0.010	0.015
$\overline{ B3 }/\overline{ B1 }$	0.086	0.028	0.006	0.056	0.035
Thick flux tube: height from reference [km]	-98	42	182	322	462
$\overline{ B2 }/\overline{ B0 }$	0.112	0.149	0.007	0.086	0.059
$\overline{ B4 }/\overline{ B0 }$	0.017	0.012	4.75 e-04	0.011	0.003
$\overline{ B3 }/\overline{ B1 }$	0.183	0.142	0.094	0.023	0.034

* Vertical bars indicate absolute values and overlines indicate horizontal average over the sheet's cross-section

panels, and less good for the two lower ones. We notice that for the three upper panels there is a systematic offset between the actual values and the thin sheet model. This is due to the fact that the sheet is slightly inclined towards the right (more positive B_x than negative in Fig. 7). This can also be seen in Fig. 4 at heights above ≈ 150 km.

B_z and B_x for the thin flux sheet can be written in a dimensional way (see also Eqns 1 and 2):

$$B_z = B_{z0} + x^2 B_{z2} + x^4 B_{z4} + \dots = B0 + B2 + B4 + \dots, \quad (16)$$

$$B_x = x B_{x1} + x^3 B_{x3} + \dots = B1 + B3 + \dots, \quad (17)$$

We can compare the relative importance of successive terms in these series expansions. Table 2 indicates that the average values $\overline{|B2|}$ are significantly smaller than $\overline{|B0|}$. The importance of $\overline{|B2|}$ is more pronounced in the upper part of the atmosphere. This is also noticeable in the upper panels of Figure 6. In a similar way to Sect. 2.2 we calculate 4th-order terms (see also Pneuman et al. (1986) and Ferriz-Mas & Schüssler (1989)). Table 2 shows that $\overline{|B4|}$ terms are very small compared to $\overline{|B0|}$. Their relative importance reaches its maximum in the top part of the atmosphere, though they remain negligible in practical terms. Similarly, $\overline{|B3|}$ terms are very much smaller than $\overline{|B1|}$. Thus the influence of successive terms in Eqs. (16) and (17) decrease with their order. This is clearly seen in Figures 6 and 7, and confirms that neglecting the 4th-order terms in B_z is justified.

5.2. Analysis of a broad flux concentration

In this section, we compare B_z and B_r from a thick flux concentration with the thin tube model (Sect. 2.1). The criteria for the choice of a flux tube in the MHD simulations are primarily its width and a relative smoothness of P_{tot} across it. The selected flux tube is located in the lower right part of Fig. 1 (crossed by a dark line). The first thing to note is that the tube is split near the




Fig. 6. Vertical component of the magnetic field, B_z , along the 5 horizontal lines in Figure 4 (solid lines). The triangles represent B_z resulting from a 2^{nd} -order thin flux sheet model.

solar surface, which probably results from the history of its interaction with convection. We also notice that this "tube" has a cross-section which deviates significantly from a circular area (see Fig. 1).

In spite of these facts, the thin flux tube model reproduces reasonably well the overall shape of B_z given in the three upper panels of Fig. 8. In the upper two, we notice the existence of a region with smoother decrease of B_z at the left edge of the flux tube. This results from a small neighbouring magnetic structure that merges with the main flux tube. This structure is not visible in the lower panels since at those heights it does not overlap the dark line (Fig. 1). It appears at the highest panels because its expansion with height makes it reach the location of the cut in the MHD cube. We don't aim to reproduce this neighbouring structure, but only the main flux tube.

In the deeper layers of the photosphere (lower panels of Fig. 8) the relatively thick flux tube splits down its center into two parts. The two separate parts of the flux tube in the lower




Fig. 7. Full lines: horizontal component of the magnetic field, B_x , across the flux sheet plotted along the 5 horizontal cuts in Fig. 4. The triangles represents B_x resulting from a 2^{nd} -order thin flux sheet model.

photosphere merge while expanding with height. It is interesting that such groups of flux concentrations tend to behave like a single flux tube higher up in the atmosphere owing to expansion and the decrease of β with height.

The splitting of the flux tube in the lowest panel leads to a decrease of the horizontally averaged field strength at this height compared to the second-lowest panel. It is seen in the framework of the thin flux tube model as an expansion of the flux tube with depth and produces positive values of h_2 (Eq. 6), clearly seen in the lowest panel.

The radial component of the magnetic field fits reasonably well with the thin flux tube model for the three upper panels (Fig. 9), except at the left edge where the small magnetic feature has merged with the main flux tube. In the two lower panels the actual profiles of B_r are disturbed by




Fig. 8. Vertical component of the magnetic field of a thick flux tube in the MHD snapshot (solid lines) along the black line in the lower right corner of Fig. 3. The triangles represent B_z resulting from the thin flux tube model. The 5 plots correspond to the same heights as in Fig. 6.

the double structure of the flux tube. In this case the thin flux tube model cannot be expected to reproduce the actual profiles. In the lowest panel, B_r from the thin flux tube model has a negative slope due to the expansion of the flux tube with depth, which leads to negative values of f_1 (Eq. 7).

The 4th-order terms remain very small at all altitudes compared to lower orders (Table 2). At the three upper altitudes, the 3rd-order contribution is clearly less marked than the 1st-order one. The uneven flux distribution at the two lower altitudes results in somewhat higher contributions of the 2nd-, 3rd- and 4th- orders compared to the situation at higher altitudes.




Fig. 9. Radial component of the magnetic field across the dark line in the lower right corner of Fig. 3. Solid lines represent B_r from the simulation, and triangles are from the thin flux tube model.

5.3. Very thin flux concentration

The thin flux tube/sheet model is generally thought to be best suited to describe the smallest flux concentrations in the MHD simulations. This picture is appropriate for the ideal case where flux tubes/sheets have an extremely thin boundary layer (separating magnetic and non-magnetic regions) and for a static plasma. The situation in the photosphere is clearly different. There vigorous convective flows induce considerable distortions of very thin flux concentrations. As a consequence, the shape and flux density distribution of the thinnest magnetic elements may differ significantly from a thin flux tube/sheet model.

In order for a flux concentration to evolve as a coherent structure in a plasma with density ρ and velocity V , its magnetic energy density ($B^2/(8\pi)$) has to be larger than the kinetic energy




Fig. 10. Vertical 2D cut in the MHD data showing B_z at the location indicated by the black line in the upper right part of the maps in Fig. 1.

density of the flow ($0.5\rho V^2$). In other words, the magnetic field has to be such that $B > B_{eq} = V\sqrt{4\pi\rho}$, where B_{eq} is the equipartition field strength.

At the surface of the sun we have $B_{eq} \simeq 500$ G. This is a limit below which we cannot expect to obtain a structure coherent enough to be described by the thin flux tube/sheet model. Thus we only consider thin magnetic features with $B > B_{eq} > 500$ G (see contours on Fig. 1). We also require that flux concentrations remain coherent at higher altitude (see top-left panel in Fig. 3) and are not located in a region close to opposite-polarity fields, since at these locations the field morphology gets complicated.

The B_z map in the upper left panel of Fig. 3 indicates that only relatively few very-thin flux concentrations (that have not merged with larger magnetic features) are noticeable at the greatest

height. We select one of them, at the location shown by the black vertical line in the upper-right part of the maps in Fig. 1.

A lateral 2D view of this thin structure (Fig. 10) shows that it is asymmetric and distorted, with an inclination that varies strongly with height. Since the magnetic energy density is not far above the equipartition value, the convective flows influence the morphology of the thin flux concentration rather strongly. This does not favour of the representation of very thin flux concentrations in terms of thin flux tube/sheet models.

6. Conclusions

The total pressure diagnostic (Sect. 4) indicates that P_{tot} is nearly constant across most flux concentrations near the solar surface. This is a necessary condition for applying the 0th-order thin flux tube/sheet approximation. In the higher parts of the atmosphere, tension forces become important due to the curved field lines and low plasma β . In this case, higher orders in the thin flux tube/sheet model are needed to describe flux elements.

For a detailed analysis of magnetic features in the MHD simulation, we have adopted two models (thin flux tube and thin flux sheet) depending on the geometry of the studied flux concentration. We have seen that for flux concentrations with magnetic field well above the equipartition distribution (Sects 5.1 and 5.2), the models reproduce reasonably well B_z and B_x (or B_r) of the simulated flux concentrations. This was especially the case in the higher part of the atmosphere. The fits were less good in the lower part of the atmosphere due to higher β and the vigorous convective flows. In this case, it is rather the overall shape of B_z that is consistent with the approximation. The 2nd-order terms of the thin flux tube/sheet approximation contribute at the 5-15 percent level especially in the upper part of the atmosphere. The 3rd-order terms provide a relatively small contribution to B_x or B_r , while the 4th-order terms give a very small contribution to B_z . This justifies neglecting the 4th-order terms and the view that higher-orders contribute less and less to B_z and B_x (or B_r).

In the case of very thin flux concentrations which generally have energy densities lower than or at most somewhat higher than the equipartition value, field lines are distorted and partly driven by plasma motions. This leads to distorted or incoherent flux concentrations which do not have the necessary symmetry and regularity to be reproduced by a thin flux tube/sheet model. To what extent these low field strengths are due to the limited resolution (low Reynold's number) of the simulations still needs to be established. Note, however, that it has been pointed out (Venkatakrisnan 1986) that the convective collapse mechanism, thought to be responsible for the concentration of magnetic flux to kG strength (Parker 1978; Spruit 1979; Grossmann-Doerth et al. 1998), becomes less efficient as the amount of magnetic flux per feature decreases. A decrease in field strength with decreasing magnetic flux has been observationally confirmed (Solanki et al. 1996).

Acknowledgements: This work was partly supported by the WCU grant No. R31-10016 from the Korean Ministry of Education, Science and Technology. L.Y.C. is thankful to the Max-Planck-Institut für Sonnensystemforschung, Katlenburg-Lindau for a stipend of the International Max Planck Research School on Physical Processes in the Solar System and Beyond.

References

- Bercik, D. J. 2002, PhD thesis, Michigan state university
- Bruls, J. H. M. J. & Solanki, S. K. 1995, *A&A*, 293, 240
- Defouw, R. J. 1976, *ApJ*, 209, 266
- Ferriz-Mas, A. & Schüssler, M. 1989, *Geophysical and Astrophysical Fluid Dynamics*, 48, 217
- Ferriz-Mas, A., Schüssler, M., & Anton, V. 1989, *A&A*, 210, 425
- Grossmann-Doerth, U., Schüssler, M., & Steiner, O. 1998, *A&A*, 337, 928
- Martínez Pillet, V., Lites, B. W., & Skumanich, A. 1997, *ApJ*, 474, 810
- Nordlund, A. 1983, in *IAU Symposium*, Vol. 102, *Solar and Stellar Magnetic Fields: Origins and Coronal Effects*, ed. J. O. Stenflo, 79–83
- Nordlund, Å. & Stein, R. F. 1990, in *IAU Symposium*, Vol. 138, *Solar Photosphere: Structure, Convection, and Magnetic Fields*, ed. J. O. Stenflo, 191
- Parker, E. N. 1978, *ApJ*, 221, 368
- Pneuman, G. W., Solanki, S. K., & Stenflo, J. O. 1986, *A&A*, 154, 231
- Rabin, D. 1992, *ApJ*, 391, 832
- Roberts, B. & Webb, A. R. 1978, *Sol. Phys.*, 56, 5
- Roberts, B. & Webb, A. R. 1979, *Sol. Phys.*, 64, 77
- Rüedi, I., Solanki, S. K., Livingston, W., & Stenflo, J. O. 1992, *A&A*, 263, 323
- Schüssler, M. 1992, in *NATO ASIC Proc. 373: The Sun: A Laboratory for Astrophysics*, ed. J. T. Schmelz & J. C. Brown, 191
- Solanki, S. K. 1993, *Space Science Reviews*, 63, 1
- Solanki, S. K., Inhester, B., & Schüssler, M. 2006, *Reports on progress in physics*, 69, 563668
- Solanki, S. K., Zufferey, D., Lin, H., Rueedi, I., & Kuhn, J. R. 1996, *A&A*, 310, L33
- Spruit, H. C. 1979, *Sol. Phys.*, 61, 363
- Spruit, H. C. 1981, *A&A*, 102, 129
- Stein, R. F. & Nordlund, A. 1998, *ApJ*, 499, 914
- Stein, R. F. & Nordlund, Å. 2003, in *Astronomical Society of the Pacific Conference Series*, Vol. 288, *Stellar Atmosphere Modeling*, ed. I. Hubeny, D. Mihalas, & K. Werner, 519
- Stenflo, J. O. 1973, *Sol. Phys.*, 32, 41
- Venkatakrishnan, P. 1986, *Nature*, 322, 156
- Vögler, A. 2003, PhD thesis, Universität Göttingen
- Vögler, A. & Schüssler, M. 2003, *Astronomische Nachrichten*, 324, 399
- Vögler, A., Shelyag, S., Schüssler, M., et al. 2005, *Astron. & Astrophys.*, 429, 335

Wiehr, E. 1978, *Mitteilungen der Astronomischen Gesellschaft Hamburg*, 43, 134

Zayer, I., Solanki, S. K., & Stenflo, J. O. 1989, *A&A*, 211, 463